













- Proximal femoral locking plate versus dynamic hip screw for unstable intertrochanteric femoral fractures. *J Orthop Surg (Hong Kong)* 2013;21:317-22.
22. Mardani-Kivi M, Mirbolook A, Khajeh Jahromi S, Rouhi Rad M. Fixation of intertrochanteric fractures: Dynamic hip screw versus locking compression plate. *Trauma Mon* 2013;18:67-70.
  23. Zhong B, Zhang Y, Zhang C, Luo CF. A comparison of proximal femoral locking compression plates with dynamic hip screws in extracapsular femoral fractures. *Orthop Traumatol Surg Res* 2014;100:663-8.
  24. Huang SG, Chen B, Zhang Y, Nie FF, Ju L, Li M, *et al.* Comparison of the Clinical Effectiveness of PFNA, PFLCP, and DHS in Treatment of Unstable Intertrochanteric Femoral Fracture. *Am J Ther* 2015:1-8.
  25. Asif N, Ahmad S, Qureshi OA, Jilani LZ, Hamesh T, Jameel T. Unstable intertrochanteric fracture fixation – Is proximal femoral locked compression plate better than dynamic hip screw. *J Clin Diagn Res* 2016;10:RC09-13.
  26. Saha P, Ayan S, Bandyopadhyay U, Mukhopadhyay AS, Bhattacharya G, Mukhopadhyay KK. Anatomical reconstruction of unstable trochanteric fractures through posterior approach. *J Orthop Allied Sci* 2015;3:55-8.
  27. Ftouh S, Morga A, Swift C; Guideline Development Group. Management of hip fracture in adults: Summary of NICE guidance. *BMJ* 2011;342:d3304.